

L8 ANSWER 4 OF 4 CAPLUS COPYRIGHT 2008 ACS on STN
 ACCESSION NUMBER: 1970:90461 CAPLUS <<LOGINID::20080430>>
 DOCUMENT NUMBER: 72:90461
 ORIGINAL REFERENCE NO.: 72:16437a,16440a
 TITLE: Anthelmintic 5-substituted aminobenzimidazoles
 INVENTOR(S): Hoff, Dale R.; Fisher, Michael Herbert
 PATENT ASSIGNEE(S): Merck and Co., Inc.
 SOURCE: S. African, 78 pp.
 CODEN: SFXXAB
 DOCUMENT TYPE: Patent
 LANGUAGE: English
 FAMILY ACC. NUM. COUNT: 1
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
ZA 6800351		19690717	ZA	
CA 967962			CA	
DE 1695545			DE	
DE 1795711			DE	
FR 1580153			FR	
GB 1198941			GB	
GB 1198942			GB	
US 3646049		19720229	US	19700305
US 3743738		19730703	US	19710602
PRIORITY APPLN. INFO.:			US	19670203
			US	19680109

GI For diagram(s), see printed CA Issue.

AB The title compds. (I) are anthelmintics, particularly useful in treating veterinary nematode infections (given orally) and also possess antifungal properties. To a solution of 10 g II (R1 = H, R2 = 4-thiazolyl) in 20 ml concentrated H2SO4 was added dropwise at 20-30° a mixture of 4 ml concentrated HNO3 and 6 ml concentrated H2SO4, and the mixture kept 5 min to give II (R1 =

O2N, R2 = 4-thiazolyl) (III), m. 240-1° (HCONMe2) (DMF). A solution of 2.2 g furfural in 3 ml EtOH was added to 3 g 3,4-(H2N)2C6H3-NO2 (IV) suspended in 10 ml PhNO2, and the mixture stirred 10 min at room temperature then heated slowly to 210° 1 min (allowing the EtOH to dist.) to give II (R1 = O2N, R2 = 2-furyl), m. 224°. II (R1 = O2N, R2 = 2-pyrryl), m. 259-60° was prepared by refluxing a mixture of 43.2 g 2-formylpyrrole, 54 g IV, and 160 g Cu(OAc)2 in 1 l. MeOH 2 hr. Catalytic hydrogenation of 141 g III in 4 l. EtOH over 22 g 5% Pd-C at 24° and 45 psi 5.5 hr gave II (R1 = H2N, R2 = 4-thiazolyl) (V), m. 232-3° (EtOH-Hexane). Addition of 1 g ClCO2Me to a suspension of 2.16 g V in 7.5 ml C5H5N gave II (R1 = MeO2CNH, R2 = 4-thiazolyl) (Va), m. 237-9° (MeOH containing C), also obtained in the absence of C5H5N using Me2CO as solvent at room temperature

Similarly were prepared the following II (R2 = 4-thiazolyl) (R1 and m.p. given): EtO2CNH, 203-5° (MeCN-Et2O) (methanolate m. 94-105°); PrO2CNH, 214-15°; BuO2CNH, 211-12°; C5H11O2CNH, 178-9°; C6H13O2CNH, 150-2°; n-C8H17O2CNH, 66-7°; PhO2CNH, 115-16°; 4-FC6H4O2CNH, 275-80°; 2-FC6H4O2CNH, 135-40°; iso-BuO2CNH, 231-2°; Me2CHO2CNH, 212-14°; CH2:CHCH2O2CNH, 210-12°; HC.tplbond.CCH2O2CNH,

200-2°; EtSc(O)NH, 215°; and cyclopropyloxycarbonylamino, 190-5° (hydrate), 207-8° (anhydrous). By standard acylation procedures were prepared the following I (R1 = H, X = O), (R2 and m.p. given): Me, 280°; PhCH2, 210-11°; H, 247-8°; Et, 255-6°; Ph (VI), 118-20°; 3-pyridyl, 284-5°; 2-FC6H4, 132-3°; 1-adamantyl, 246-7°; 2-naphthyl, 154-6°; cyclopropyl, 245°; Me2CH, 203-5°; 3-thienyl, 276-8°; 3-FC6H4, 232-3°; 4-FC6H4, 151-2°; 2-MeOC6H4, 113-14°; 3-MeOC6H4, 105-9°; 2-PhO, 95-100°; 2-ClC6H4, 146-7°; 3-IC6H4, 127-9°; 3-FC6H4, 201-3°; 3-ONC6H4, 163-4°; 2,5-F2C6H3, 113-14°; 2-pyridyl, 240-1°; 4-pyridyl, 150-3°; Me3C, 241-2°; 2-furyl, 139-40°; 4-thiazolyl, 387-8°; 2-thienyl, 288° (decomposition); MeOCH2, 238-9°; Cl2CH, 220°; and Me2C:CH, 270-2°. Refluxing a mixture of 5 g I (R1 = H, R2 = EtS, X = O), (VIa) 0.5 g Bu2SnO, and 50 ml F3CCH2OH 20 hr gave I (R1 = H, R2 = F3CCH2O), m. 231-2° (AcOEt-hexane). I (R1 = H, R2 = HC.tplbond.CCH2O, X = O), m. 202-2°, was similarly prepared. Addition of 3.62 g PhOC(S)Cl dropwise to 4.32 g V in 25 ml C5H5N and stirring 1.5 hr gave I (R1 = H, R2 = PhO, X = S), m. 155-7° (MeOH), which on heating in C5H5N (7 vols.) 1 hr at 100° gave 5-isothiocyanato-2-(4-thiazolyl)benzimidazole (VII), m. 243-6°. I (R1 = H, R2 = MeO, X = S), m. 224° (MeOH), was prepared by refluxing a mixture of 2.5 g VII, 25 mg NaOMe, and 300 ml MeOH 20 hr; I (R1 = H, R2 = EtO, X = S), prepared similarly, m. 218°. A slow stream of MeSH was passed into 4 g VII in 25 ml Me2NCHO 15 min, and the mixture kept 20 hr to give I (R1 = H, R2 = MeS, X = S), m. 202-5°. I (R1 = H, R2 = Ph, X = S), m. 140-3° (MeOH), was obtained by refluxing a mixture of 1 g VI, 2 g P2S5, and 20 ml C5H5N 25 min. A suspension of V (amount unspecified) in 25 ml C5H5N was treated dropwise with 1.2 g MeNCO and the mixture stirred 2.5 hr to give I (R1 = H, R2 = MeNH, X = O), m. 160° (MeOH). Dropwise addition of 2.5 g Me2NCOCl to a suspension of 4.32 g V in 25 ml C5H5N gave I (R1 = H, R2 = Me2N, X = O), m. 260-2° (MeOH). Refluxing a mixture of 5 g VIa and 25 ml Et2NH 6 hr gave I (R1 = H, R2 = Et2N, X = O), m. 234-5°. The following I (R1 = H) were similarly prepared (R2, X, and m.p. given): pyrrolidino, O, 296-8°; piperidino, O, -; MeNH, S, 235-7°; PhNH, S, 244-6°; Me2N, S, 156-9°; Et2N, S, 130-5°; piperidino, S, 225-6°, and pyrrolidino, S, 257-8°. A mixture of 8.5 g I (R1 = H, R2 = Me2CHO, X = O) (VIII), 100 ml DMF and 2.3 g 52% NaOH emulsion in mineral oil was stirred at room temperature 20 min., heated 10 min at 50°, cooled to room temperature, treated with 7.1 g MeI in 10 ml DMF, and the mixture heated 20 min at 80° to give I (R1 = Me, R2 = Me2CHO, X = O), purified by crystallization from AcOEt. Addition of a mixture of 1.3 ml concentrated HNO3 (d. 1.41) and 2.8 ml concentrated H2SO4 to 3.8 g 1-methoxy-2-(4-thiazolyl)benzimidazole in 12.3 ml concentrated H2SO4 at 12° gave 1.5 g 5-nitro-1-methoxy-2-(4-thiazolyl)benzimidazole (IX), m. 220-1° (MeOH). Hydrogenation of IX in MeOH and treatment with methanolic HCl gave 450 mg amorphous 5-amino analog-HCl, which with ClCO2CHMe2 in C5H5N gave I (R1 = MeO, R2 = Me2CHO, X = O), m. 123-5°. Treatment of 5.4 g Va in 100 ml PhMe and 30 ml DMF with 0.7 g NaH in 2 ml PhMe at 65°, stirring 1 hr, adding 2.5 g AcCl at 55°, and refluxing 0.5 hr gave I (R1 = Ac, R2 = MeO, X = O). A mixture of 3.26 g VIII and 1 g BuNCO in 100 ml MeCN was refluxed 4 hr to give I (R1 = BuNHCO, R2 =


Me₂CHO, X = O). Treatment of VIII in C₅H₅N with ClCO₂CHMe₂ gave I (R₁ = Me₂CHO₂C, R₂ = Me₂CHO, X = O), crystallized from aqueous MeOH. Similarly, V and MeSO₂Cl gave II (R₁ = MeSO₂NH, R₂ = 4-thiazolyl), m. 225-6° (MeOH). II (R₁ = PhSO₂NH, R₂ = 4-thiazolyl) (X), m. 254-5°, was also prepared. Methylation of 3.5 g X in 10 ml MeOH containing 0.54 g NaOMe with 0.625 ml MeI gave II (R₁ PhSO₂NMe, R₂ = 4-thiazolyl), m. 142-3° (MeOH); II (R₁ = MeSO₂NMe, R₂ = 4-thiazolyl), m. 192-3° (MeCN), was also prepared. The following XI were analogously prepared (R₁, R₂, R₃, and m.p. given): H, MeO₂CNMe, 4-thiazolyl, 161-2°; H, MeO₂CNH, 2-furyl, 162-3°; MeO, MeO₂CNH, 2-furyl, 164°; H, EtO₂CNH, 2-furyl, 171-2°; H, PhO₂CNH, 2-furyl, 150-5°; H, EtO₂CNH, 2-pyrrol, 200-2°; H, MeO₂CNH, 2-thienyl, 185-8°; H, MeO₂CNH, 1,2,5-thiadiazol-3-yl, 150-5°; H, MeO₂CNH, 1-pyrazolyl, 207-10°; H, MeO₂CNH, 2-methyl-4-thiazolyl, 135°; H, MeO₂CNH, 1,2,3-thiadiazol-4-yl, 218-20°; H, MeO₂CNH, 1,3,4-thiadiazol-2-yl, 258°; H, Me₂CHO₂CNH, 2-oxazolyl, 206°; H, Me₂CHO₂CNH, 2-thiazolyl, 234°; H, MeO₂CNH, 2-imidazolyl, 205-7°; H, 4-FC₆H₄CONH, 2-furyl, 264°; H, 2-furoylamino, 2-furyl, 248°; H, 4-FC₆H₄CONH, 1-pyrazolyl, 230°; H, BzNH, 2-thiazolyl, 135-40°.

IT 27217-07-8P 27217-08-9P 27217-09-0P
28767-12-6P

RL: SPN (Synthetic preparation); PREP (Preparation)
(preparation of)


RN 27217-07-8 CAPLUS

CN Urea, N,N-dimethyl-N'-[2-(4-thiazolyl)-1H-benzimidazol-5-yl]- (9CI) (CA INDEX NAME)




RN 27217-08-9 CAPLUS

CN Urea, 1,1-diethyl-3-[2-(4-thiazolyl)-5-benzimidazolyl]- (8CI) (CA INDEX NAME)



RN 27217-09-0 CAPLUS

CN 1-Pyrrolidinecarboxamide, N-[2-(4-thiazolyl)-5-benzimidazolyl]- (8CI) (CA INDEX NAME)



RN 28767-12-6 CAPLUS

CN Urea, 1-methyl-3-[2-(4-thiazolyl)-5-benzimidazolyl]- (8CI) (CA INDEX NAME)

